





Tekturna
(aliskiren) tablets 150 mg-300 mg



Tekturna HCT
(aliskiren and hydrochlorothiazide) tablets
150 mg/12.5 mg • 150 mg/25 mg • 300 mg/12.5 mg • 300 mg/25 mg

Savings Card

**Pay As Little As
\$40 For Your
30-Day Prescription***

*Reimbursement limited to \$40 per prescription. Additional limitations apply.

OpusHealth™

RxBIN: 601341
RxPCN: OHCP
RxGrp: OH3101051
RxID: C61100764852
Suf: 01

Patient Instructions: Present this card to your pharmacist, along with your insurance card (if applicable) and a valid prescription for TEKTURNA® or TEKTURNA HCT®, to receive up to a \$40 savings on your prescription co-pay after you pay the first \$40. This offer is valid for up to 12 30-day fills of TEKTURNA or TEKTURNA HCT. Cardholders with questions, please call 1-800-471-9342 (8:30am – 5:30pm ET, Monday – Friday).

Pharmacist Instructions: This card must be accompanied by a valid prescription for TEKTURNA or TEKTURNA HCT. Please submit the copay authorized by the patient's primary insurance as a secondary transaction to Opus Health. For self-pay patients, submit the claim at U&C. Pharmacists with questions, please call Opus Health at 1-800-364-4767 (8:30am – 5:30pm ET, Monday – Friday and 8:30am – 2pm ET, Saturday).

This card is the property of Noden Pharma USA Inc. and Opus Health and must be returned upon request. Not valid in any state where prohibited by law. Both parties retain the right to rescind, revoke, or amend this program at any time without notice.

Visit www.tekturna.com for full Prescribing Information and Patient Information, including BOXED WARNING, Contraindications, and Warnings and Precautions.

Offer Valid For Insured And Cash Patients

1. This offer is not valid if prescriptions are paid by any state or other federally funded programs, including, but not limited to, Medicare, Medicaid, Medigap, VA, DoD, or TRICARE®.
2. This card is good for use with a valid TEKTURNA or TEKTURNA HCT prescription at the time the prescription is filled by the pharmacist and dispensed to the patient.
3. Program is only valid for patients aged 18 years and older.
4. The selling, purchasing, trading, or counterfeiting of this card is prohibited by law.
5. By using this card, you demonstrate that you understand and agree to comply with the terms and conditions of this offer as set forth on this card.
6. Card is limited to one per person and is not transferable.
7. This card is NOT health insurance. Actual payment for product will depend on individual insurance coverage.

Please see the Important safety Information on the following page.



IMPORTANT SAFETY INFORMATION

IMPORTANT WARNING: TEKTURNA (aliskiren) or TEKTURNA HCT (aliskiren and hydrochlorothiazide) can cause harm or death to an unborn baby. Talk to your doctor about other ways to lower your blood pressure if you plan to become pregnant. If you become pregnant, stop taking TEKTURNA or TEKTURNA HCT and tell your doctor right away.

Administer or take TEKTURNA or TEKTURNA HCT as instructed by your Doctor. TEKTURNA should not be given to children under 2 years of age. The safety and effectiveness of TEKTURNA has not been established in pediatric patients younger than 6 years.

Do not take TEKTURNA or TEKTURNA HCT if you have diabetes and are taking a kind of medicine called an angiotensin receptor blocker (ARB) or angiotensin-converting enzyme inhibitor (ACEI); or if you are allergic to aliskiren or any of their other ingredients.

In addition, do not take TEKTURNA HCT if you make very little or no urine due to kidney problems, or if you have allergic reactions to certain medicines called sulfonamides or hydrochlorothiazide.

Aliskiren, a medicine in TEKTURNA and TEKTURNA HCT, can cause difficulty breathing or swallowing, tightness of the chest, hives, general rash, swelling, itching, dizziness, vomiting, or abdominal pain (signs of a severe allergic reaction called angioedema). TEKTURNA and TEKTURNA HCT can also cause swelling of the face, lips, tongue, throat, arms and legs, or the whole body (signs of angioedema). Stop taking TEKTURNA or TEKTURNA HCT, get medical help right away and tell your doctor if you get any one or more of these symptoms. Angioedema can happen at any time while you are taking TEKTURNA or TEKTURNA HCT.

Your blood pressure may get too low if you also:

- take water pills (also called “diuretics”)
- are on a low-salt diet
- get dialysis treatments
- have heart problems
- get sick with vomiting or diarrhea
- drink alcohol and take certain medicines (barbiturates or narcotics). These can cause low blood pressure to get worse with TEKTURNA HCT.

Lie down if you feel faint or dizzy. Call your doctor or seek medical attention right away.

Tell your doctor about all of your medical conditions, including: if you have kidney problems; if you have ever had an allergic reaction, called angioedema, to another blood pressure medicine; or if you are breastfeeding. It is not known if TEKTURNA or TEKTURNA HCT passes into your breast milk and can harm your baby. You should not take TEKTURNA or TEKTURNA HCT and breastfeed.

Tell your doctor right away if you have eye pain or a decrease in vision.

Especially, tell your doctor about all the medicines you take, including a kind of medicine to control blood pressure, called an angiotensin receptor blocker (ARB) or ACE inhibitor (ACEI), cholesterol lowering medicines (like simvastatin, atorvastatin, cholestyramine, colestipol), water pills (especially potassium-sparing diuretics), medicines to treat fungal infections (like itraconazole or ketoconazole), cyclosporine, potassium-containing medicines, potassium supplements, salt substitutes containing potassium, or NSAIDs (like ibuprofen or naproxen), including Cyclooxygenase-2 inhibitors (COX-2 inhibitors), and non-prescription medicines, vitamins and herbal supplements.

Additionally, if you take TEKTURNA HCT, tell your doctor if you have liver problems, have allergies or asthma, have lupus, or if you take lithium or medicine to treat diabetes.

The most common side effects of TEKTURNA and TEKTURNA HCT include:

- diarrhea
- cough
- dizziness
- headache
- flu-like symptoms
- back pain
- tiredness
- high levels of potassium in the blood (hyperkalemia)
- vertigo
- joint pain (arthralgia)

Less common side effects include skin rash, severe skin reactions, liver disorder, and low level of sodium in blood.

Store this medicine at room temperature between 68°F to 77°F (20°C to 25°C) and keep out of reach of children. Keep in its original container and protect from moisture.

You are encouraged to report side effects of prescription drugs to the FDA.

To report SUSPECTED ADVERSE REACTIONS, contact Noden Pharma USA Inc. at 1-844-399-5701 and/or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

Please see full Prescribing Information and Patient Information, including **IMPORTANT WARNING**, for TEKTURNA.

Please see full Prescribing Information and Patient Information, including **IMPORTANT WARNING**, for TEKTURNA HCT.